



Toxic elements in blood of red-necked nightjars (*Caprimulgus ruficollis*) inhabiting differently polluted environments[☆]



Silvia Espín ^{a, b, *}, Pablo Sánchez-Virosta ^{a, b}, José M. Zamora-Marín ^{c, d},
Mario León-Ortega ^e, Pedro Jiménez ^a, Manuel Zumbado ^{f, g}, Octavio P. Luzardo ^{f, g},
Tapiro Eeva ^b, Antonio J. García-Fernández ^a

^a Toxicology and Risk Assessment Group, Department of Health Sciences, IMIB-Arrixaca, Faculty of Veterinary, University of Murcia, Campus de Espinardo, 30100, Murcia, Spain

^b Department of Biology, University of Turku, Turku, Finland

^c Department of Zoology and Physical Anthropology, Faculty of Biology, University of Murcia, Murcia, Spain

^d Grupo de Anillamiento ANSE, Murcia, Spain

^e Department of Ecology and Hydrology, Faculty of Biology, University of Murcia, Murcia, Spain

^f Toxicology Unit, Research Institute of Biomedical and Health Sciences, University of Las Palmas de Gran Canaria, Las Palmas, Spain

^g Spanish Biomedical Research Center in Physiopathology of Obesity and Nutrition (CIBERObn), Spain

ARTICLE INFO

Article history:

Received 7 November 2019

Received in revised form

3 March 2020

Accepted 4 March 2020

Available online 5 March 2020

Keywords:

Metal exposure

Nightjars

Caprimulgus ruficollis

Industrial emissions

Urbanization

ABSTRACT

Toxic metals have been widely reported in avian tissues due to their well-known accumulation capacity and adverse effects. However, rare earth elements (REE) and other minor elements (ME) are becoming a new threat due to their use in modern technology. Presently, exposure data are limited and no studies have been reported in wildlife. The order Caprimulgiformes is among the most understudied groups of birds lacking blood ecotoxicological data. One major constraint is the small blood sample volume that can be collected to ensure animal welfare, which limits analyses. In order to shed light on these collective issues, we sampled 48 red-necked nightjars (*Caprimulgus ruficollis*) inhabiting three different scenarios of contaminant exposure (agricultural-urban area, n = 15; mining area, n = 17; and control area, n = 16) in southeastern Spain, and report for the first time concentrations of 50 elements (i.e. trace elements, ATSDR's list toxic elements, REE and ME) using a recently developed technique able to analyze them by ICP-MS in very small volumes of blood (130 µL). Concentrations of As, Cd, Pb and Mn were significantly higher in individuals captured at the mining area compared to the other sites. Lead levels in the mine site were of particular concern since it was in the range of blood concentrations related to subclinical/clinical effects in other species, and in our study were associated with decreased hematocrit values (up to 44% hematocrit depression at blood concentrations >1000 ng/ml w.w.). Moreover, additive effects related to metal cocktail exposure in the mining area could be expected. Age and gender-related differences in blood concentrations were found for some elements. Even though most REE and ME concentrations were close to the LOQ, some of these emerging contaminants may trigger sublethal effects that, together with the ATSDR's list toxic elements, need to be carefully evaluated in a future study.

© 2020 Elsevier Ltd. All rights reserved.

1. Introduction

Toxic elements have been widely reported in avian tissues due to their well-known bioaccumulation capacity and the adverse effects they cause in birds, being able to alter reproductive success and behavior, compromise immune function and affect different biochemical parameters (Eeva et al., 2005; Espín et al., 2016a, 2016b; Pain et al., 2019; Sánchez-Virosta et al., 2015; Vallverdú-Coll et al., 2019; Whitney and Cristol, 2018). However, the scientific

* This paper has been recommended for acceptance by Wen-Xiong Wang.

* Corresponding authors. Area of Toxicology, Faculty of Veterinary, University of Murcia, Campus de Espinardo, 30100, Murcia, Spain.

E-mail addresses: silvia.espin@um.es (S. Espín), pablo.s.v@um.es (P. Sánchez-Virosta), josemanuel.zamora@um.es (J.M. Zamora-Marín), mlovertega@um.es (M. León-Ortega), pjjm@um.es (P. Jiménez), manuel.zumbado@ulpgc.es (M. Zumbado), octavio.perez@ulpgc.es (O.P. Luzardo), teeva@utu.fi (T. Eeva), ajgf@um.es (A.J. García-Fernández).

community agrees that further research evaluating metal exposure is required to assess spatial and temporal trends and fill the knowledge gap for some avian species, and even for entire orders of birds from which no data exist. In this way, biomonitoring studies can provide early warning of potential impacts in wildlife species and the wider environment, and can be used to track the success of mitigation in reducing exposure (Espín et al., 2016c; García-Fernández, 2014).

Some metalloids and metals such as arsenic (As), lead (Pb), mercury (Hg) or cadmium (Cd) are clearly considered toxic and are ranked in the Substance Priority List (1st, 2nd, 3rd and 7th position, respectively) prepared by the Agency of Toxic Substances and Disease Registry (ATSDR, 2017). Rare earth elements (REE) and other minor elements (ME) are becoming increasingly important due to their intensive use in modern technology all over the world, generating aerial emissions and tons of electronic waste (Hussain and Mumtaz, 2014; Tansel, 2017). However, data on exposure to these elements have been rarely reported (e.g. in humans: Gaman et al., 2019; González-Antuña et al., 2017), with no studies in wildlife. Thus, there is much uncertainty about their accumulation capacity and the potential short and long-term consequences for wild avian populations. This may be of particular concern to those species migrating to developing/under developed countries receiving hazardous waste from more developed countries and where almost all the e-waste recycling chain reaches the informal sector (collection, manual dismantling, open burning to recover metals and open dumping of residuals), with the consequent impact on health and the environment (Tansel, 2017; UNEP, 2011).

Nightjars (*Caprimulgiformes*) are crepuscular and nocturnal insectivorous birds widespread over the world, including more than 120 species distributed on all continents except in arctic climates and New Zealand (Cleere, 2010). Despite wide geographic distribution, nightjars are among the most poorly studied groups of birds, since their nocturnal habits, cryptic plumage and elusive behavior have traditionally hampered the study of their biology and ecology (Braun and Huddleston, 2009). Recently, information for a few species has increased due to the development of suitable monitoring and sampling methods (e.g. Camacho et al., 2016; Evens et al., 2017), but many general aspects of their relation with human activities and responses to habitat degradation remain still unknown. For example, to our knowledge, no studies have been conducted to explore the exposure and effects of environmental pollutants in any nightjar species, despite of the variety of pollutants emitted worldwide from a range of sources such as agriculture, industry and mining activities.

The Red-necked nightjar (*Caprimulgus ruficollis*, hereafter called "nightjar") is a long-distance migrant quite common in warm regions of the Iberian Peninsula and northern Africa, where it breeds from early spring to late summer (Aragonés, 2003; Camacho, 2013a). Until few years ago, this species has been considered one of the least known birds in its distribution area (Aragonés, 2003; Sáez-Gómez et al., 2015). Wintering grounds are located on Western Sahara, presumably in countries such as Mali, Guinea Bissau and Senegal, although the exact range is unclear (Cleere et al., 2013). From dusk to dawn, nightjars use open shrublands, crops and roads to hunt aerial insects through short flights or taking prey directly from the ground (Camacho, 2013b; Jackson, 2003). Main prey species of nightjars are nocturnal insects belonging to orders Lepidoptera, Homoptera and Neuroptera (Camacho, 2013a), although its diet has not been thoroughly explored. In Spain, field observations have reported moths (Noctuidae and Thaumetopoeidae families, among others) as the main species composing the nightjar diet (Sáez and Camacho, 2016), however, other insect

families such as Gryllidae and Cicadidae can contribute significantly to nightjar diet when their populations peak (authors' unpublished data). Nightjars show a notable tolerance to anthropogenic alterations, being able to reach high population densities even in highly managed landscapes (Camacho et al., 2014). Due to this ecological plasticity, nightjars can occur from natural Mediterranean shrublands to mining-impacted areas or intensive irrigation crops. However, toxic elements exposure and potential effects from industrial activities (i.e. mining exploitation or intensive agriculture) on nightjar physiology and fitness condition have never been studied. Hence, this study aims to evaluate the exposure to 50 elements (i.e. trace elements, ATSDR's list toxic elements, REE and ME) in nightjars inhabiting three different scenarios of contaminant exposure in the southeast of Iberian Peninsula. We expect increased Pb concentrations in the mining area based on previous findings in other species, showing δALAD inhibition (Espín et al., 2015) that may lead to pathologic effects including anemia, reduced hematocrit and hemoglobin or even brain damage (Dieter and Finley, 1979; Hoffman et al., 2000). Thus, potential effects of Pb exposure on hematocrit levels are also tested. Finally, we also evaluate the influence of factors at individual level, such as age and sex, on the observed concentrations of these elements in whole blood.

2. Material and methods

2.1. Study area

This study was conducted in the province of Murcia, in the southeast of Iberian Peninsula (37° 45'N, 0° 57'W) (Fig. 1). The study area is characterized by a Mediterranean semi-arid climate, with a strong water deficit during spring and summer and scarce rainfall occurring predominantly in winter. Main land use of the study area is irrigation agriculture, which extends over lowlands, while highland landscapes are dominated by a mosaic of rainfed crops, pine forests and Mediterranean shrublands.

Three different scenarios of contaminant exposure were selected in the study area according to the main land use in each one: agricultural-urban area, mining area and control area. Agricultural-urban area is located at the center of the province of Murcia, next to Guadalentín River, and is occupied by large extensions of lucerne and cotton crops adjacent to an industrial park and a densely populated city. Contamination sources in this area are not known, but it is expected that intense agriculture and industrial activities lead to moderate pollution levels. Mining area corresponds to an ancient mining site (Cartagena-La Unión Mining District) located on the south of the province of Murcia, with extraction activity since Phoenicians, Carthaginians and Roman times until the end of the 20th century (ceased in 1992) (Conesa et al., 2008). Significant pollutant levels have been found in blood samples of wildlife in this area, mainly for metals such as Pb, Hg and Cd, among others (Espin et al., 2014a, 2014b; García-Fernández et al., 1995). Furthermore, toxic metals are still spread by small creeks from headwaters, due to the eroding process of runoff waters, impacting on surrounding ecosystems (Conesa and Schulz, 2010). Finally, we selected Escalona and Altaona mountains as a control area, where no contaminants are known to affect wildlife (Espin et al., 2014b). Human occupancy in this area is considerably low and landscape is dominated mainly by small tree crops (citric, almond and olive trees) scattered in a shrubland and forest matrix (León-Ortega et al., 2017). Then, pollution sources are scarce and localized in this area (Espin et al., 2014a, 2014b).




Fig. 1. Map showing the geographical location of the studied areas.

2.2. Sampling and measurements

In June and July 2017, seven sampling nights were conducted in the three different pollution scenarios, and a total of 48 nightjars were captured ($n = 15$ at agricultural-urban area, $n = 17$ at mining area and $n = 16$ at control area). From dusk to dawn, we drove a car at constant speed (15–20 km/h) over secondary roads distributed throughout the study areas. Nightjars were detected by the eye shine triggered when *tapetum lucidum* was lighting on frontal view. When detected, birds were captured using a torch and handheld net with 10 mm mesh-size (Camacho et al., 2014; Jackson, 2003). Once captured, each nightjar was individually marked with metal ring, and the age, sex, weight and other biometric measures (not used in this study) were recorded. Age and sex ($n = 19$ males and $n = 29$ females) were determined by feather design in wing and tail, following available literature for this species (Alonso and Caballero, 2003; Forero et al., 1995; Gargallo, 1994; Tornero and Sanchís, 2017). We grouped birds into three age classes: juveniles (young birds in their first summer – EURING age code III, $n = 5$), breeding subadults (birds in their second summer – EURING age code V, $n = 8$) or breeding adults (older birds – EURING age code VI, $n = 35$).

The clinical examination of each bird was performed by a veterinarian prior to blood sampling. No symptoms were observed in the individuals sampled, being considered clinically healthy. Blood samples (approximately 1 mL) were collected by puncturing brachial veins with 30G needles and 1 mL-syringes and stored in heparinized Eppendorf tubes under refrigerated conditions until processed in the laboratory. Hematocrit (% of red blood cells from total sample volume) was recorded using capillary tube readers after centrifugation of blood at 2200 g for 5 min. One Eppendorf tube with whole blood was frozen at -80°C until element analysis. Other Eppendorf tube with whole blood was centrifuged to separate plasma and red blood cell, and the tubes were frozen at -80°C for biochemical analysis that will be the scope of a different publication. Duration of the handling process per sampled nightjar ranged from 10 to 15 min and all birds were released exactly in the same location where they were caught.

2.3. Trace element analysis

We determined whole blood concentrations of 50 elements (Table 1), which were selected according to their toxicity and/or their frequent use in the manufacture of electronic consumer products (Hussain and Mumtaz, 2014; Tansel, 2017). For the elemental analysis we employed an Agilent 7900 ICP-MS equipment (Agilent Technologies, Tokyo, Japan) equipped with standard nickel cones, Ultra High Matrix Introduction (UHMI) system, and a cross-flow nebulizer with a Make-Up Gas Port (X400 Nebulizer, Savillex Corporation, MN, USA). We followed a procedure developed for human blood, which had been previously validated in our laboratory using certified reference materials (González-Antuña et al., 2017). Briefly, 130 μL of blood were diluted using 1120 μL of ammonia solution (0.05% of EDTA, 0.05% of Triton X-100, and 1% of NH_4OH), and 50 μL of internal standards (ISTD) were added (final volume = 1.3 mL). ISTD solution consisted of Sc (scandium), Ge (germanium), Rh (rhodium), and Ir (iridium) at a stock concentration of 20 mg/mL each. Pure standards of elements in acid solution (5% HNO_3 , 100 mg/L) were purchased from CPA Chem (Stara Zagora, Bulgaria). Two standard curves (ten points, 20 ng/mL – 0.005 ng/mL) were made to avoid interferences between elements: a) one using a commercial multi-element mixture (CPA Chem, 100 mg/L, 5% HNO_3) containing all the essential elements and main heavy metals; and b) other multi-element mixture tailor-made in our laboratory from individual elements (CPA Chem), which contained the REE and ME. The limits of quantification (LOQs) ranged between 0.005 and 1.0 ng/ml, and the accuracy of measurements were in the range of 79–138%, with relative standard deviations (RSD) below 6% in all cases, as previously described (González-Antuña et al., 2017).

2.4. Statistical procedures

For all 50 elements we first calculated ranges and quartiles (Table 1). Most of the elements showed some values below the limit of quantification (<LOQ) and for many of them this proportion was relatively high (Table 1). Therefore, for statistical comparison we only selected those elements where the proportion of <LOQ values was below 15% (EPA, 2000). For these 17 elements we

Table 1

Range and quartiles (Q25, median, Q75) for element concentrations (ng/ml, w.w.) in whole blood of the red-necked nightjar (*Caprimulgus ruficollis*). Lower limit of quantification (LOQ) given together with the proportion of samples < LOQ, N = 48.

Element	Group ^a	Range (min – max)	Q25	Median	Q75	LOQ	%<LOQ
Aluminum (Al)	2	<LOQ – 1300	11.3	23.8	29.6	38.4	88
Antimony (Sb)	2	<LOQ – 3.22	0.010	0.010	0.010	0.010	88
Arsenic (As)	2	0.250–58.3	0.800	2.26	6.82	0.008	0
Barium (Ba)	2	<LOQ – 235	16.9	25.9	37.1	1.02	4
Beryllium (Be)	2	<LOQ – 0.356	0.020	0.039	0.045	0.050	81
Bismuth (Bi)	4	<LOQ – 0.233	0.006	0.007	0.008	0.050	98
Cadmium (Cd)	2	<LOQ – 5.35	0.235	0.450	1.46	0.010	10
Cerium (Ce)	3	<LOQ – 4.84	0.608	0.873	1.38	0.050	6
Chromium (Cr)	1	<LOQ – 14.1	1.06	1.71	3.78	0.229	6
Cobalt (Co)	1	<LOQ – 10.1	0.702	1.10	2.53	0.011	4
Copper (Cu)	1	152–477	284	352	395	1.72	0
Dysprosium (Dy)	3	<LOQ – 0.064	0.008	0.010	0.023	0.005	52
Erbium (Er)	3	N/A	<LOQ	<LOQ	<LOQ	0.050	100
Europium (Eu)	3	<LOQ – 0.021	0.007	0.008	0.010	0.005	46
Cadolinium (Gd)	3	<LOQ – 0.119	0.009	0.026	0.037	0.005	29
Gallium (Ga)	4	<LOQ – 0.917	0.167	0.232	0.296	0.050	6
Gold (Au)	4	<LOQ – 0.756	0.171	0.414	0.523	0.005	69
Holmium (Ho)	3	<LOQ – 0.010	0.006	0.007	0.008	0.005	79
Indium (In)	4	<LOQ – 0.028	0.007	0.007	0.011	0.005	69
Iron (Fe)	1	80800–657000	422000	470000	522000	24.6	0
Lanthanum (La)	3	<LOQ – 4.99	0.040	0.088	0.232	0.100	56
Lead (Pb)	2	<LOQ – 1320	10.1	16.9	182	0.361	4
Lutetium (Lu)	3	<LOQ – 0.010	0.006	0.007	0.009	0.005	96
Manganese (Mn)	1	5.73–327	13.5	18.5	32.8	0.371	0
Mercury (Hg)	2	5.83–351	15.6	26.0	38.3	0.010	0
Molybdenum (Mo)	1	11.7–44.1	24.6	30.5	37.2	0.148	0
Neodymium (Nd)	3	<LOQ – 0.471	0.007	0.009	0.190	0.005	58
Nickel (Ni)	1	<LOQ – 206	2.46	4.52	6.43	7.95	94
Niobium (Nb)	4	<LOQ – 0.317	0.008	0.065	0.097	0.005	38
Osmium (Os)	4	N/A	<LOQ	<LOQ	<LOQ	0.005	100
Palladium (Pd)	2	N/A	<LOQ	<LOQ	<LOQ	0.010	100
Platinum (Pt)	4	<LOQ – 0.010	0.006	0.008	0.009	0.005	98
Praseodymium (Pr)	3	<LOQ – 0.125	0.007	0.008	0.010	0.005	81
Ruthenium (Ru)	4	<LOQ – 0.025	0.007	0.008	0.009	0.005	83
Samarium (Sm)	3	<LOQ – 0.094	0.006	0.008	0.009	0.005	88
Selenium (Se)	1	145–1880	694	943	1180	0.153	0
Silver (Ag)	2	<LOQ – 12.8	0.031	0.073	0.088	0.100	79
Strontium (Sr)	2	14.0–665	37.4	70.1	116	0.439	0
Tantalum (Ta)	4	<LOQ – 0.104	0.007	0.010	0.014	0.005	10
Terbium (Tb)	3	<LOQ – 0.012	0.006	0.008	0.009	0.005	85
Thallium (Tl)	2	<LOQ – 6.74	0.029	0.041	0.059	0.050	73
Thorium (Th)	2	<LOQ – 0.135	0.024	0.032	0.046	0.050	88
Thulium (Tm)	3	<LOQ – 0.010	0.006	0.007	0.009	0.005	98
Tin (Sn)	2	<LOQ – 4.32	0.010	0.010	0.510	0.010	56
Titanium (Ti)	4	<LOQ – 56.4	14.9	21.5	25.0	0.757	23
Uranium (U)	2	<LOQ – 0.050	0.023	0.027	0.042	0.050	98
Vanadium (V)	2	<LOQ – 6.39	0.027	0.763	1.07	0.050	35
Ytterbium (Yb)	3	<LOQ – 0.031	0.006	0.008	0.010	0.005	48
Yttrium (Y)	3	<LOQ – 0.279	0.009	0.070	0.118	0.005	38
Zinc (Zn)	1	2810–19200	4860	5680	7030	51.0	0

^a Element categories: 1 = Essential trace elements, 2 = ATSDR's list toxic elements, 3 = Rare earth elements, 4 = Other minor elements.

substituted < LOQ values by a random number between 0 and LOQ. Three of the elements (Cu, Fe, Mo) were normally distributed (by visual inspection of histogram and Kolmogorov-Smirnov test for normality) while the rest of the elements were log₁₀ transformed to make them better conform normal distribution. After that, we analyzed differences among sampling areas by linear models (LM) with area, sex and age as explanatory factors. Because we found relatively high Pb values in nightjar blood we further tested the association between Pb level and hematocrit with a generalized linear model (GLM; using beta error distribution and logit link function) where Pb level was used as an explanatory factor together with sex and age). Model estimates (least squares means and confidence limits) for log₁₀-transformed values were back-transformed to the original scale for tables and figures. For the elements that showed significant differences among areas or age classes we further ran Tukey's test adjusted for the number of

pairwise comparisons. Alpha level was set to 0.05 in all analyses. Finally, associations among elements were inspected with a hierarchical cluster analysis by using Pearson correlation matrix with average linkage method for clustering. All the analyses were ran with a statistical software SAS 9.4 ([SAS Institute Inc, 2013](#)).

3. Results and discussion

Many of the elements analyzed (33 out of 50, mainly REE and ME) were below LOQ in more than 15% of the individuals ([Table 1](#)), most often indicating low general levels and suggesting limited effects at the population level. For those elements where the proportion of < LOQ values was below 15%, mean concentrations in whole blood of nightjars by sampling area and GLMs are shown in [Table 2](#). In general, element concentrations did not differ among the areas. However, concentrations of As, Cd, Pb and Mn were

Table 2

Mean (\pm 95% confidence limits) element concentrations (ng/ml, w.w.) in whole blood of the red-necked nightjar (*Caprimulgus ruficollis*) at three sampling environments (agricultural-urban area, mining area, control area). Linear models (LM) for comparison of means. Tukey's test: means with the same letter are not statistically different. N = 48.

Element	Mean			LM ^a <i>F</i> _{2,42}	<i>p</i>
	Agricultural-Urban (CL)	Mining (CL)	Control (CL)		
Arsenic (As)*	1.17 ^a (0.658–2.09)	6.87 ^b (3.56–13.2)	1.56 ^a (0.907–2.69)	13.0	<0.0001
Barium (Ba)*	23.0 (10.9–48.3)	29.8 (12.8–69.2)	17.0 (8.45–34.2)	0.72	0.49
Cadmium (Cd)*	0.093 ^a (0.042–0.208)	0.609 ^b (0.244–1.52)	0.119 ^a (0.056–0.254)	7.80	0.0013
Cerium (Ce)*	0.943 (0.450–1.98)	0.610 (0.263–1.41)	0.625 (0.312–1.25)	0.54	0.58
Chromium (Cr)*	1.82 (0.945–3.52)	1.66 (0.789–3.51)	1.45 (0.783–2.70)	0.16	0.86
Cobalt (Co)*	0.753 (0.344–1.65)	1.53 (0.627–3.72)	1.33 (0.634–2.77)	1.12	0.34
Copper (Cu)	294 (245–344)	289 (233–345)	323 (277–370)	0.74	0.49
Gallium (Ga)*	0.224 (0.147–0.342)	0.179 (0.111–0.289)	0.181 (0.121–0.269)	0.45	0.64
Iron (Fe)	483000 (421000–544000)	434000 (372000–496000)	431000 (377000–485000)	0.67	0.52
Lead (Pb)*	7.65 ^a (3.63–16.2)	166 ^b (71.0–386)	3.87 ^a (1.92–7.81)	34.5	<0.0001
Manganese (Mn)*	13.6 ^a (8.15–22.5)	30.5 ^b (17.2–54.3)	23.8 ^{ab} (14.8–38.5)	3.28	0.048
Mercury (Hg)*	15.4 (9.92–23.8)	21.2 (12.9–34.7)	24.8 (16.4–37.4)	1.59	0.22
Molybdenum (Mo)	31.2 ^{ab} (27.0–35.4)	25.1 ^a (20.3–29.9)	33.4 ^b (29.4–37.4)	5.19	0.0097
Selenium (Se)*	878 (679–1140)	680 (508–909)	880 (691–1120)	1.64	0.21
Strontium (Sr)*	51.1 (33.1–79.1)	62.6 (38.2–103)	63.9 (42.4–96.3)	0.40	0.67
Tantalum (Ta)*	0.009 (0.006–0.015)	0.013 (0.008–0.022)	0.009 (0.006–0.015)	0.81	0.45
Zinc (Zn)*	5180 (4180–6420)	6020 (4720–7680)	5360 (4380–6560)	0.65	0.53

*Geometric means, values log₁₀ transformed for the analyses and back-transformed for the table.

^a Explanatory factors in the models were study area, age and sex.

significantly increased in blood of individuals captured at the mining area compared to the other sites (for Mn only compared to the agricultural-urban area), while Mo concentrations were reduced in nightjars from the mining area compared to the control site (Table 2).

Birds are widely used in biomonitoring studies to evaluate toxic element contamination from industrial activities, and insectivorous birds have proved to be useful to assess metal exposure because they accumulate metals through the diet (Berglund et al., 2011; Eeva et al., 2009; Espín et al., 2016d; García-Fernández, 2014; Sánchez-Virosta et al., 2015). In our study, the diet is considered the main exposure route of toxic metals in nightjars (i.e. ingestion of prey that have accumulated the elements), as reported for other species (Espin et al., 2014a, 2014b) and in general in different bird studies (Espín et al., 2016c; García-Fernández et al., 1995; Outridge and Scheuhammer, 1993; Sánchez-Virosta et al., 2015). Field observations have reported that the diet of nightjars is mainly composed by nocturnal aerial insects, with moths representing the main prey species in Spain (Sáez and Camacho, 2016). In this sense, larvae of moths (Lepidoptera) have shown increased toxic element concentrations (e.g. As, Cd, Pb) in metal-polluted environments compared to reference sites (Eeva et al., 2018). However, as explained before, the diet of nightjars has not been thoroughly described due to methodological constraints. Therefore, further studies are needed to explore the diet composition, as well as to develop suitable sampling methods for metal analyses in prey items.

Once absorbed, elements are transported and distributed throughout the organism via the blood, and their concentrations in internal tissues are a key indicator of bioaccumulation (Espín et al., 2016c). The main accumulation organ depends on the toxicokinetics of each element and element form, as well as the type of exposure: e.g. bone is the main organ of Pb accumulation but there is also distribution to liver, kidney and brain; kidney and liver are the main organs for Cd and Hg accumulation; liver is the main organ for As (Dauwe et al., 2002; Espín et al., 2016c; García-Fernández et al., 1995, 1996, 1997; Sánchez-Virosta et al., 2015, 2018). The main excretion rate and route also depends on the element and species physiology, being urine and feces the main route, although some metals can be sequestered in feathers during molt or in the eggs/eggshells during laying (Espín et al., 2012,

2016c; Martínez-López et al., 2004, 2005; Sánchez-Virosta et al., 2015). Overall, blood levels provide a non-destructive measure of contaminant exposure (Espín et al., 2016c). Blood is a proxy of short-term exposure to metals, and although it could reflect part of the metals accumulated in wintering grounds (for adults birds) and mobilized during breeding, it mostly reflects recent exposure in the breeding area (Espín et al., 2016c). Due to the lack of ecotoxicological data not only in the Caprimulgidae family, but in the whole Caprimulgiformes taxa, element concentrations found in this study were compared to those reported in different wild birds (mainly feeding on insects when possible). In this study, Pb is of particular concern due to the marked differences between the mining area and the other sites, nightjars in the mine site showing Pb concentrations ca. 43 × higher than in the control area. Previous studies developed during the 1990s and 2010s demonstrate that birds inhabiting close to this ancient mine site show higher Pb concentrations (Espin et al., 2014b, 2015; García-Fernández et al., 1995) due to the intensive mining activity (mainly of Pb, Zn, Cu, Mn, Fe, Ag and Sn) developed for more than 2500 years until its closing in 1992 (Conesa et al., 2008; Pavetti et al., 2006), being the principal Pb and Zn source in Spain during the 19th century (Estevan-Senís, 1967). Blood Pb concentrations found in nightjars inhabiting the mining area (arithmetic mean = 443 ng/ml w.w.) were similar to those found in red-winged blackbirds (*Agelaius phoeniceus*) at contaminated sites in USA (Tsiopoura et al., 2008) (Fig. 2). However, in general, Pb concentrations in nightjars were higher than those reported in American robins (*Turdus migratorius*), feral pigeon (*Columba livia*), northern mockingbird (*Mimus polyglottos*), common blackbirds (*Turdus merula*), little owl (*Athene noctua*), common kestrel (*Falco tinnunculus*) and gray catbird (*Dumetella carolinensis*) inhabiting polluted environments in USA, the Netherlands, France and Spain, the latter species exhibiting worse body condition as a result of Pb contamination (Cai and Calisi, 2016; García-Fernández et al., 1995, 1997; Roux and Marra, 2007; Scheifler et al., 2006; Schilderman et al., 1997) (Fig. 2).

Regarding other elements, As blood concentrations in the mining area were similar to those found in red-winged blackbirds at polluted sites in USA (Tsiopoura et al., 2008), and Cd levels were lower than those reported in blackbirds, and similar to those found in common kestrel, little owl (García-Fernández et al., 1995), and nestling eagle owls (*Bubo bubo*) in the same mine site (Espin et al.,




Fig. 2. Blood Pb concentrations (ng/ml, w.w.) found in different bird species inhabiting polluted/urban areas in the literature.

2014b). These As and Cd concentrations are within the range considered as low exposure levels in birds (Espin et al., 2014b; García-Fernández et al., 1995; Tsipoura et al., 2008). Both Mn and Mo are essential elements, but exposure to high levels may cause adverse effects (Stafford et al., 2016; Williams et al., 2012). Blood concentrations found in nightjars in the mining area were within the range of values found by other researchers in apparently healthy birds: Mo levels were similar to those reported in urban areas in France for great tit (*Parus major*) (19 ng/ml) (Bailly et al., 2017) and lower than those found in Northwestern crows (*Corvus caurinus*) in USA (55–66 ng/ml w.w.) (Van Hemert and Handel, 2016), while Mn concentrations were similar to those found in Northwestern crows (28–33 ng/ml w.w.) and much lower than those reported in African white-backed vultures (*Pseudogyps africanus*) in South Africa (162–196 ng/ml w.w. in juveniles and 473 ng/ml w.w. in adults) (Van Hemert and Handel, 2016; van Wyk et al., 2001).

Pb levels in the mining area were in the range of blood concentrations related to subclinical signs in birds, 65% of individuals exceeding 200 ng/ml w.w. (considered the benchmark value for subclinical and physiological effects for Anseriformes and Falconiformes), 35% exceeding 500 ng/ml (considered clinical poisoning) and 2 individuals (12%) exceeding 1000 ng/ml in blood (related to severe clinical poisoning in Anseriformes and Falconiformes) (Franson and Pain, 2011). A previous study in the same mining area showed that nestling eagle owls exposed to Pb suffer up to 79% δALAD inhibition at blood Pb concentrations ≥190 ng/ml (Espín et al., 2015). Some researchers suggest that high inhibitions (>75%) for this enzyme may lead to pathologic effects including anemia, reduced hematocrit and hemoglobin or even brain damage (Dieter and Finley, 1979; Hoffman et al., 2000). We found that hematocrit of red-necked nightjars was negatively associated to blood Pb level (GLM: $F_{1,43} = 10.4$, $p = 0.0024$; estimate = -0.00052), but did not depend on age ($F_{2,43} = 1.0$, $p = 0.37$) or sex ($F_{1,43} = 1.6$, $p = 0.21$). Hematocrit values decreased by 4, 12 and 44% (hematocrit: 37, 34 and 22%) at blood concentrations ranging 200–500, 500–1000 and exceeding 1000 ng/ml, respectively, as compared with the mean hematocrit value (38%) in nightjars from the less Pb-polluted areas, with blood Pb concentrations averaging 12.8 ng/ml (maximum of 31.6 ng/ml). Hematocrit is a marker of general health (Boross et al., 2012), and its depression in some individuals of this

study, possibly partly due to Pb exposure, may be indicative of deteriorated health condition. One study showed that a 94% decrease in δALAD activity was needed to cause a decline of 11% in hematocrit in griffon vulture (*Gyps fulvus*) (Espín et al., 2015), suggesting that nightjars showing high Pb concentrations could present an important δALAD enzyme inhibition. In addition, previous studies in the mining area showed that blood concentrations as low as 30 ng/ml for Pb or 0.2 ng/ml for Cd were able to deplete major antioxidants of the red blood cells (i.e. glutathione, glutathione peroxidase, catalase) and induce lipid damage in nestling eagle owls (Espín et al., 2014b). This suggests that the classical threshold level for Pb (and possibly for other elements) commonly accepted for considering physiological effects in certain avian species should be revised in the wild, where individuals are exposed to Pb but also to a combination of other contaminants and stressors. Therefore, potential sublethal effects of some elements, mainly Pb but also Cd, As and others, on nightjar's physiology cannot be excluded, and we consider that they will most likely occur. Moreover, additive effects related to metal cocktail exposure in the mining area could be expected.

Fallon et al. (2017) published some guidelines for evaluation of Pb poisoning of wild raptors, presenting the criteria for removing a Pb-poisoned bird from the wild for treatment: (i) if blood Pb concentration is low (≤ 400 ng/ml) and the bird appears healthy, it should be returned to the wild, (ii) if blood Pb concentration is elevated (400–600 ng/ml) but clinical signs are not present, the bird should be returned to the wild, but if clinical signs are present, the bird should be transferred to a rehabilitator for treatment, and (iii) if blood Pb concentrations are toxic (≥ 600 ng/ml), the bird should be transferred to the rehabilitator regardless of the symptoms. Chelation therapy is the main treatment for Pb poisoning, using chelation agents such as D-penicillamine, British Anti-lewisite (BAL), dimercaptosuccinic acid (succimer) and CaNa₂ ethylene diamine tetra acetic acid (CaEDTA), the two later being considered the treatments of choice (Redig and Arent, 2008). Fallon et al. (2017) inform that these categories should be extrapolated with caution to different raptor species. Although the blood Pb levels of nightjars were below those thresholds in most individuals, it must be recalled that the clinical effects vary among species: e.g. clinical signs in swans (*Cygnus sp.*) have been reported at ≥ 400 ng/ml and lethal values are > 2000 ng/ml (Degernes et al., 2002; Sears

et al., 1989). Therefore, further studies are needed to understand how different species respond to Pb exposure (Fallon et al., 2017). This issue should be carefully evaluated in a future study to ascertain how nightjars' physiology and condition respond to Pb exposure.

Sublethal effects of Pb exposure on breeding performance should also be considered. Pb-related impacts on avian reproduction have been well documented (e.g. reduced fertilization, hatching rate, breeding success, sperm quality) (Fritsch et al., 2019; Gasparik et al., 2012; Vallverdú-Coll et al., 2016). Vallverdú-Coll et al. (2016) experimentally exposed red-legged partridges (*Alectoris rufa*) to Pb shot pellets, and one month after exposure females and males showed blood Pb levels of 557 ± 29 and 821 ± 149 ng/ml respectively in the low dose group, and 758 ± 75 and 920 ± 202 ng/ml in the high dose group. These concentrations were related to lower hatching and reduced acrosome integrity and sperm motility (Vallverdú-Coll et al., 2016). Some nightjars in the mining area showed equivalent blood Pb concentrations. Efforts to monitor breeding and ecological parameters in nightjars in the study area have just started, thus reproductive data will be gathered in next years to evaluate the population trend and the potential reproductive consequences of Pb exposure.

Four elements (Cu, Mn, Mo and Sr) showed sex differences, females (all of them sampled in breeding season) showing higher concentrations than males (Fig. 3). This species does not show sexual dimorphism in body size or body mass (Camacho et al., 2016; Forero et al., 1995), so this would not explain the gender-related differences found here. Cu, Mn and Mo are essential elements required during the egg development, which need mineral stores deposited in the egg at the time of its formation for the proper embryogenesis (Miles, 2000; Vieira, 2007). Thus, gender-related differences in the physiological state, with breeding females showing higher levels of some essential elements to be transferred to the egg, could explain this result. In addition, food intake of laying females could also be higher, increasing accumulation from the environment, or females in their reproductive period could choose prey with greater nutritional value (proteins, lipids, calcium, other essential elements, etc.), as occurs in other bird species. On the other hand, two elements (Cd, Hg) showed age-specific differences, with the lowest concentrations in the youngest age class (Fig. 4). Increased concentrations in adults may be explained by the longer time to bioaccumulate and increase their




Fig. 4. Means ($\pm 95\%$ confidence limits) of two elements showing age-specific differences in blood concentrations of the red-necked nightjar (*Caprimulgus ruficollis*). N = 48 (for age classes: III = 5, V = 8, VI = 35). Tukey's test: means with the same letter (within element) are not statistically different.

metal body burden (Evers, 2018; Sánchez-Virosta et al., 2015; Wayland and Scheuhammer, 2011). In this sense, accumulation by age seems to be very typical especially for Cd, even in wild animals exposed to low background levels of dietary Cd (Berglund et al., 2011; Scheuhammer, 1987). In addition, although blood concentrations mainly reflect recent exposure (primarily via ingestion) (Burger et al., 2018; Espín et al., 2016c), they also partly reflect bioaccumulated elements that transfer from tissues to blood (Burger et al., 2018). Thus, since juveniles birds are in their first summer and they have not spent the winter in Africa yet, it is possible that their subadult/adult counterparts have accumulated higher levels of those specific metals in their wintering grounds (located on Western Sahara, (Cleere et al., 2013)). The greater effectiveness in finding food for adults compared to young birds, also increases the possibility of contaminant exposure in older birds. Finally, potential age or gender-related differences in diet could also explain these results. However, no studies on age or




Fig. 3. Means ($\pm 95\%$ confidence limits) of four elements showing sex-specific differences in blood concentrations of the red-necked nightjar (*Caprimulgus ruficollis*). N = 48 (19 males + 29 females).




Fig. 5. Hierarchical clustering of 17 elements in whole blood of the red-necked nightjar (*Caprimulgus ruficollis*) on the basis of their Pearson correlation matrix. N = 48 for each element.

gender-related differences in feeding behavior and diet composition are available for this species.

The toxic elements Pb, Cd, As and Hg were grouped in a cluster analysis (Fig. 5), indicating correlation among them and suggesting common origin in the polluted site. Correlations among the less toxic elements might reflect homeostatic regulation which controls absorption and hence body trace element concentrations (e.g. essential elements like Co, Mn and Zn, or Se and Fe). However, the clusters generated could also be explained by other reasons such as diet composition, element similarities in atomic structure/reactivity, and soil composition (physicochemical factors may control element associations during soil formation, e.g. Co and Mn are closely associated in soils because their similar chemical properties, Bradford et al., 1996; Uren, 2013).

4. Conclusions

Due to the lack of ecotoxicological studies on Caprimulgiformes taxa, this article reports for the first time blood concentrations of 50 elements (i.e. trace elements, ATSDR's list toxic elements, REE and ME) in red-necked nightjars (*Caprimulgus ruficollis*) inhabiting three different scenarios of contaminant exposure, using a recently developed ICP-MS technique requiring small sample volumes. Blood As, Cd, Pb and Mn concentrations were significantly increased in nightjars captured at the mining area compared to the control and agricultural-urban areas. Lead levels in the mine site are of particular concern since they were in the range of blood concentrations related to subclinical and clinical effects in others avian taxa, and in our study were associated with decreased hematocrit values (up to 44% hematocrit depression at blood concentrations exceeding 1000 ng/ml w.w.). A few elements (Cu, Mn, Mo and Sr) showed higher values in females than in males, and two elements (Cd and Hg) accumulated with the age. It is well known that metals and metalloids such as Pb, Cd, Hg or As may produce oxidative stress and alter calcium metabolism. Thus, sublethal effects of some elements (mainly Pb) on nightjar's physiology with potential medium and long-term consequences cannot be discarded. Moreover, additive effects related to metal cocktail exposure in the mining area could be expected. Even though most REE and ME concentrations were close to the LOQ, some of these emerging contaminants may trigger sublethal effects that, together with the ATSDR's list toxic elements, needs to be evaluated in a future study.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

CRediT authorship contribution statement

Silvia Espín: Conceptualization, Methodology, Formal analysis, Writing - original draft, Funding acquisition. **Pablo Sánchez-Virosta:** Methodology, Writing - review & editing. **José M. Zamora-Marín:** Methodology, Writing - review & editing. **Mario León-Ortega:** Methodology, Writing - review & editing. **Pedro Jiménez:** Methodology, Writing - review & editing. **Manuel Zumbado:** Methodology, Writing - review & editing. **Octavio P. Luzardo:** Methodology, Writing - review & editing. **Tapio Eeva:** Formal analysis, Writing - review & editing. **Antonio J. García-Fernández:** Methodology, Writing - review & editing.

Acknowledgment

This work was supported by the Fundación Séneca - Agencia de Ciencia y Tecnología de la Región de Murcia (20031/SF/16 to S.E.). S.E. was financially supported by Fundación Séneca (*Saavedra-Fajardo* Project 20031/SF/16) and by Ministerio de Ciencia, Innovación y Universidades (*Juan de la Cierva-Incorporación* contract, IJCI-2017-34653). We thank the COST Action 16224 ERBFa-cility for maintaining an international collaboration network for researchers. We also thank two anonymous reviewers for their useful comments and suggestions.

All applicable institutional and/or national guidelines for the care and use of animals were followed.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envpol.2020.114334>.

References

- Alonso, R., Caballero, M.J., 2003. Determinación de la edad y sexo en *Caprimulgus ruficollis*. Revista de Anillamiento 17–22.
- Aragonés, J., 2003. Breeding biology of the Red-necked Nightjar *Caprimulgus ruficollis* in southern Spain. ARDEOLA 50, 215–221.
- ATSDR, 2017. The ATSDR 2017 Substance Priority List. U.S.A.: Agency for Toxic Substances and Disease Registry, U.S. Public Health Service.
- Bailly, J., Faivre, B., Bernard, N., Sage, M., Crini, N., Driget, V., Garnier, S., Rieffel, D., Scheifler, R., 2017. Multi-element analysis of blood samples in a passerine species: excesses and deficiencies of trace elements in an urbanization study. Front. Ecol. Evol. 5, 1–9.
- Berglund, Å.M.M., Koivula, M.J., Eeva, T., 2011. Species- and age-related variation in metal exposure and accumulation of two passerine bird species. Environ. Pollut. 159, 2368–2374.
- Boross, N., Markó, G., Laczi, M., Garamszegi, Z., Hegyi, G., Herényi, M., Kiss, D., Nagy, G., Rosival, R., Szöllősi, E., et al., 2012. Sources of variation in haematocrit in the collared flycatcher (*Ficedula albicollis*). Ornis Hung. 20, 64–72.
- Bradford, G., Chang, A., Page, A., Bakthar, D., Frampton, J., Wright, H., 1996. Background Concentration of Trace and Major Elements in California Soils. Kearney Foundation Special Report, University of California. Kearney Foundation, pp. 1–32.
- Braun, M.J., Huddleston, C.J., 2009. A molecular phylogenetic survey of caprimulgiform nightbirds illustrates the utility of non-coding sequences. Mol. Phylogenet. Evol. 53, 948–960.
- Burger, J., Mizrahi, D., Tsipoura, N., Jeitner, C., Gochfeld, M., 2018. Mercury, lead, cadmium, cobalt, arsenic and selenium in the blood of semipalmed sandpipers (*Calidris pusilla*) from Suriname, south America: age-related differences in wintering site and comparisons with a stopover site in New Jersey, USA. Toxics 6, 27.
- Cai, F., Calisi, R.M., 2016. Seasons and neighborhoods of high lead toxicity in New York City: the feral pigeon as a bioindicator. Chemosphere 161, 274–279.
- Camacho, C., 2013a. Tropical phenology in temperate regions. Condor 115, 830–837.
- Camacho, C., 2013b. Behavioural thermoregulation in man-made habitats: surface choice and mortality risk in red-necked nightjars. Hous. Theor. Soc. 60, 124–130.
- Camacho, C., Palacios, S., Sáez, P., Sánchez, S., Potti, J., 2014. Human-induced changes in landscape configuration influence individual movement routines: lessons from a versatile, highly mobile species. PLoS One 9 (8), e104974.
- Camacho, C., Sáez, P., Sánchez, S., Palacios, S., Molina, C., Potti, J., 2016. The road to opportunities: landscape change promotes body-size divergence in a highly mobile species. Curr. Zool. 62, 7–14.
- Cleere, N., 2010. Nightjars of the World: Potoos, Frogmouths, Oilbird and Owl-Nightjars. Ltd. WILD, Old Basing, UK.
- Cleere, N., Kirwan, G.M., Christie, D.A., de Juana, E., 2013. Red-necked nightjar (*Caprimulgus ruficollis*). In: Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona, Spain.
- Conesa, H.M., Schulin, R., 2010. The Cartagena–La Unión mining district (SE Spain): a review of environmental problems and emerging phytoremediation solutions after fifteen years research. J. Environ. Monit. 1225–1233.
- Conesa, H., Schulin, R., Nowack, B., 2008. Mining landscape: a cultural tourist opportunity or an environmental problem?: the study case of the Cartagena–La Unión Mining District (SE Spain). Ecol. Econ. 64, 690–700.
- Dauwe, T., Bervoets, L., Blust, R., Eens, M., 2002. Tissue levels of lead in experimentally exposed zebra finches (*Taeniopygia guttata*) with particular attention on the use of feathers as biomonitor. Arch. Environ. Contam. Toxicol. 42, 88–92.
- Degernes, L., Kittelson, S., Linck, M., Manthey, P., 2002. Survival and reproductive success of trumpeter swans after treatment for lead poisoning. Waterbirds: Int.

- J. Waterbird Biol. 25, 368–374.
- Dieter, M.P., Finley, M.T., 1979. δ -aminolevulinic acid dehydratase enzyme activity in blood, brain, and liver of lead-dosed ducks. Environ. Res. 19, 127–135.
- Eeva, T., Hasselquist, D., Lange fors, Å., Tummeleht, L., Nikinmaa, M., Ilmonen, P., 2005. Pollution related effects in immune function and stress in a free-living population of pied flycatcher *Ficedula hypoleuca*. J. Avian Biol. 36, 405–412.
- Eeva, T., Ahola, M., Lehikoinen, E., 2009. Breeding performance of blue tits (*Cyanistes caeruleus*) and great tits (*Parus major*) in a heavy metal polluted area. Environ. Pollut. 157, 3126–3131.
- Eeva, T., Holmström, H., Espín, S., Sánchez-Virosta, P., Klemola, T., 2018. Leaves, berries and herbivorous larvae of bilberry *Vaccinium myrtillus* as sources of metals in food chains at a Cu-Ni smelter site. Chemosphere 210, 859–866.
- EPA, 2000. Guidance for Data Quality Assessment: Practical Methods for Data Analysis (EPA QA/G9: QA00 Update: July 2000).
- Espín, S., Martínez-López, E., Gómez-Ramírez, P., María-Mojica, P., García-Fernández, A.J., 2012. Razorbills (*Alca torda*) as bioindicators of mercury pollution in the southwestern Mediterranean. Mar. Pollut. Bull. 64 (11), 2461–2470.
- Espín, S., Martínez-López, E., León-Ortega, M., Calvo, J.F., García-Fernández, A.J., 2014a. Factors that influence mercury concentrations in nestling Eagle Owls (*Bubo bubo*). Sci. Total Environ. 470–471, 1132–1139.
- Espín, S., Martínez-López, E., Jiménez, P., María-Mojica, P., García-Fernández, A.J., 2015. Delta-aminolevulinic acid dehydratase (δ ALAD) activity in four free-living bird species exposed to different levels of lead under natural conditions. Environ. Res. 137C, 185–198.
- Espín, S., Ruiz, S., Sánchez-Virosta, P., Salminen, J.-P., Eeva, T., 2016a. Effects of experimental calcium availability and anthropogenic metal pollution on eggshell characteristics and yolk carotenoid and vitamin levels in two passerine birds. Chemosphere 151, 189–201.
- Espín, S., Martínez-López, E., Jiménez, P., María-Mojica, P., García-Fernández, A.J., 2016b. Interspecific differences in the antioxidant capacity of two Laridae species exposed to metals. Environ. Res. 147, 115–124.
- Espín, S., García-Fernández, A.J., Herzke, D., Shore, R.F., van Hattum, B., Martínez López, E., Coeurdassier, M., Eulaers, I., Fritsch, C., Gómez-Ramírez, P., et al., 2016c. Tracking pan-continental trends in environmental contamination using sentinel raptors — what types of samples should we use? Ecotoxicology 25, 777–801.
- Espín, S., Martínez-López, E., León-Ortega, M., Martínez, J.E., García-Fernández, A.J., 2014b. Oxidative stress biomarkers in Eurasian eagle owls (*Bubo bubo*) in three different scenarios of heavy metal exposure. Environ. Res. 131, 134–144.
- Espín, S., Ruiz, S., Sánchez-Virosta, P., Eeva, T., 2016d. Effects of calcium supplementation on growth and biochemistry in two passerine species breeding in a Ca-poor and metal-polluted area. Environ. Sci. Pollut. Res. Int. 23, 9809–9821.
- Estevan-Senís, M.T., 1967. Explotación minera de la Sierra de Cartagena (1840–1919). Saitabi 17, 211–234.
- Evens, R., Conway, G.J., Henderson, I.G., Cresswell, B., Jiguet, F., Moussy, C., Sénacl, D., Witters, N., Beernaerts, N., Artois, T., 2017. Migratory pathways, stopover zones and wintering destinations of Western European Nightjars *Caprimulgus europaeus*. Ibis 159, 680–686.
- Evers, D., 2018. The effects of methylmercury on wildlife: a comprehensive review and approach for interpretation. In: The Encyclopedia of the Anthropocene. Elsevier, Oxford, pp. 181–194.
- Fallon, J.A., Redig, P., Miller, T.A., Lanzone, M., Katzner, T., 2017. Guidelines for evaluation and treatment of lead poisoning of wild raptors. Wildl. Soc. Bull. 41, 205–211.
- Forero, M.G., Tella, J.L., García, L., 1995. Age related evolution of sexual dimorphism in the red-necked nightjar *Caprimulgus ruficollis*. J. Ornithol. 447–451.
- Franson, J.C., Pain, D.J., 2011. Lead in birds. In: Environmental Contaminants in Biota: Interpreting Tissue Concentrations. Florida: CRC Press, Taylor & Francis, Boca Raton, pp. 563–593.
- Fritsch, C., Jankowiak, Ł., Wysocki, D., 2019. Exposure to Pb impairs breeding success and is associated with longer lifespan in urban European blackbirds. Sci. Rep. 9.
- Gaman, L., Delia, C.E., Lizardo, O.P., Zumbado, M., Badea, M., Stoian, I., Gilca, M., Boada, L.D., Henríquez-Hernández, L.A., 2019. Serum concentration of toxic metals and rare earth elements in children and adolescent. Int. J. Environ. Health Res. 1–17.
- García-Fernández, A.J., 2014. Ecotoxicology, Avian. Encyclopedia of Toxicology, 3rd edition. Elsevier Inc., Academic Press, pp. 289–294.
- García-Fernández, A.J., Sánchez-García, J.A., Jiménez-Montalbán, P., Luna, A., 1995. Lead and cadmium in wild birds in southeastern Spain. Environ. Toxicol. Chem. 14, 2049–2058.
- García-Fernández, A.J., Sánchez-García, J.A., Gómez-Zapata, M., Luna, A., 1996. Distribution of cadmium in blood and tissues of wild birds. Arch. Environ. Contam. Toxicol. 30, 252–258.
- García-Fernández, A.J., Motas-Guzmán, M., Navas, I., María-Mojica, P., Luna, A., Sánchez-García, J.A., 1997. Environmental exposure and distribution of lead in four species of raptors in Southeastern Spain. Arch. Environ. Contam. Toxicol. 33, 76–82.
- Gargallo, G., 1994. Flight feather moult in the Red-necked Nightjar *Caprimulgus ruficollis*. J. Avian 25, 119–124.
- Gasparik, J., Venglarčík, J., Slamecka, J., Kropil, R., Smehyl, P., Kopecký, J., 2012. Distribution of lead in selected organs and its effect on reproduction parameters of pheasants (*Phasianus colchicus*) after an experimental per oral administration. J. Environ. Sci. Health A Toxicol. Hazard Subst. Environ. Eng. 47, 1267–1271.
- González-Antuña, A., Camacho, M., Henríquez-Hernández, L.A., Boada, L.D., Almeida-González, M., Zumbado, M., Lizardo, O.P., 2017. Simultaneous quantification of 49 elements associated to e-waste in human blood by ICP-MS for routine analysis. MethodsX 4, 328–334.
- Hoffman, D.J., Heinz, G.H., Sileo, L., Audet, D.J., Campbell, J.K., LeCaptain, L.J., 2000. Developmental toxicity of lead-contaminated sediment to mallard ducklings. Arch. Environ. Contam. Toxicol. 39, 221–232.
- Hussain, M., Mumtaz, S., 2014. E-waste: impacts, issues and management strategies. Rev. Environ. Health 29, 53–58.
- Jackson, H.D., 2003. A field survey to investigate why nightjars frequent roads at night. Ostrich 74, 97–101.
- León-Ortega, M., Martínez, J.E., Pérez, E., Lacalle, J.A., Calvo, J.F., 2017. The contribution of non-protected areas to the conservation of Eurasian Eagle-owls in Mediterranean ecosystems. Ecosphere 8, e01952.
- Martínez-López, E., Martínez, J.E., María-Mojica, P., Peñalver, J., Pulido, M., Calvo, J.F., García-Fernández, A.J., 2004. Lead in feathers and delta-aminolevulinic acid dehydratase activity in three raptor species from an unpolluted Mediterranean forest (Southeastern Spain). Arch. Environ. Contam. Toxicol. 47, 270–275.
- Martínez-López, E., María-Mojica, P., Martínez, J.E., Calvo, J.F., Romero, D., García-Fernández, A.J., 2005. Cadmium in feathers of adults and blood of nestlings of three raptor species from a nonpolluted mediterranean forest, southeastern Spain. Bull. Environ. Contam. Toxicol. 74, 477–484.
- Miles, R.D., 2000. Trace minerals and avian embryo development. Ciencia Anim. Bras. 2, 1–10.
- Outridge, P.M., Scheuhammer, A.M., 1993. Bioaccumulation and toxicology of nickel: implications for wild mammals and birds. Environ. Rev. 1, 172–197.
- Pain, D.J., Mateo, R., Green, R.E., 2019. Effects of lead from ammunition on birds and other wildlife: a review and update. Ambio 48, 935–953.
- Pavetti, F.G., Conesa-Alcaraz, M., Faz-Cano, A., Arnaldos, R., García, G., 2006. Contaminación por metales pesados en la rambla de El Gorguel (SE Península Ibérica). Terra 24, 171–178.
- Redig, P.T., Arent, L.R., 2008. Raptor toxicology. Veterinary Clinics: Exotic Animal Pract. 11, 261–282.
- Roux, K., Marra, P., 2007. The presence and impact of environmental lead in passerine birds along an urban to rural land use gradient. Arch. Environ. Contam. Toxicol. 53, 261–268.
- Sáez, P., Camacho, C., 2016. Chotacabras Cuellirrojo — *Caprimulgus Ruficollis*. Museo Nacional de Ciencias Naturales, Madrid. <http://www.vertebradosibericos.org>.
- Sáez-Gómez, P., Camacho, C., Palacios, S., Sánchez-Navarro, S., Ruiz-Ramos, J., Molina, C., Dávila, C., Potti, J., 2015. El chotacabras cuellirrojo en Doñana: un elemento conciliador entre agricultura y conservación. Quercus 355, 25–32.
- Sánchez-Virosta, P., Espín, S., García-Fernández, A.J., Eeva, T., 2015. A review on exposure and effects of arsenic in passerine birds. Sci. Total Environ. 512 (513), 506–525.
- Sánchez-Virosta, P., Espín, S., Ruiz, S., Salminen, J.-P., García-Fernández, A.J., Eeva, T., 2018. Experimental manipulation of dietary arsenic levels in great tit nestlings: accumulation pattern and effects on growth, survival and plasma biochemistry. Environ. Pollut. 233, 764–773.
- SAS Institute Inc, 2013. Base SAS 9.4 Procedures Guide: Statistical Procedures.
- Scheifler, R., Coeurdassier, M., Morilhat, C., Bernard, N., Faivre, B., Flicoteaux, P., Giraudoux, P., Noël, M., Piotte, P., Rieffel, D., et al., 2006. Lead concentrations in feathers and blood of common blackbirds (*Turdus merula*) and in earthworms inhabiting unpolluted and moderately polluted urban areas. Sci. Total Environ. 371, 197–205.
- Scheuhammer, A.M., 1987. The chronic toxicity of aluminium, cadmium, mercury, and lead in birds: a review. Environ. Pollut. 46, 263–295.
- Schilderman, P.A.E.L., Hoogewerff, J.A., van Schooten, F.-J., Maas, L.M., Moonen, E.J.C., van Os, B.J.H., van Wijnen, J.H., Kleinjans, J.C.S., 1997. Possible relevance of pigeons as an indicator species for monitoring air pollution. Environ. Health Perspect. 105, 322–330.
- Sears, J., Cooke, S.W., Cooke, Z.R., Heron, T.J., 1989. A method for the treatment of lead poisoning in the mute swan (*Cygnus olor*) and its long-term success. Br. Vet. J. 145, 586–595.
- Stafford, J.M., Lambert, C.E., Zyskowski, J.A., Engfehr, C.L., Fletcher, O.J., Clark, S.L., Tiwary, A., Gulde, C.M., Sample, B.E., 2016. Dietary toxicity of soluble and insoluble molybdenum to northern bobwhite quail (*Colinus virginianus*). Eco-toxicology 25, 291–301.
- Tansel, B., 2017. From electronic consumer products to e-wastes: global outlook, waste quantities, recycling challenges. Environ. Int. 98, 35–45.
- Tornero, J.A., Sanchís, M.J., 2017. Patrón de muda y dato del chotacabras cuellirrojo *Caprimulgus ruficollis*. Revista de Anillamiento 36, 16–30.
- Tsipoura, N., Burger, J., Feltes, R., Yacobucci, J., Mizrahi, D., Jeitner, C., Gochfeld, M., 2008. Metal concentrations in three species of passerine birds breeding in the Hackensack Meadowlands of New Jersey. Environ. Res. 107, 218–228.
- UNEP, 2011. Where Are WEEE in Africa? Findings from the Basel Convention E-Waste Africa Programme.
- Uren, N., 2013. Cobalt and manganese. In: Heavy Metals in Soils. Environmental Pollution, Springer, Dordrecht.
- Vallverdú-Coll, N., Mougeot, F., Ortiz-Santaliestra, M.E., Castaño, C., Santiago-Moreno, J., Mateo, R., 2016. Effects of lead exposure on sperm quality and reproductive success in an avian model. Environ. Sci. Technol. 50, 12484–12492.
- Vallverdú-Coll, N., Mateo, R., Mougeot, F., Ortiz-Santaliestra, M.E., 2019. Immunotoxic effects of lead on birds. Sci. Total Environ. 689, 505–515.
- Van Hemert, C., Handel, C., 2016. Elements in whole blood of northwestern crows

- (*Corvus caurinus*) in Alaska, USA: No evidence for an association with beak deformities. *J. Wildl. Dis.* 52, 713–718.
- van Wyk, E., van der Bank, F., Verdoorn, G.H., Hofmann, D., 2001. Selected mineral and heavy metal concentrations in blood and tissues of vultures in different regions of South Africa. *S. Afr. J. Anim. Sci.* 31, 57–63.
- Vieira, S.L., 2007. Chicken embryo utilization of egg micronutrients. *Braz. J. Poultry Sci.* 9, 1–8.
- Wayland, M., Scheuhammer, A.M., 2011. Cadmium in birds. In: *Environmental Contaminants in Biota: Interpreting Tissue Concentrations*. CRC Press, Taylor & Francis, Boca Raton, pp. 645–669.
- Whitney, M.C., Cristol, D.A., 2018. Impacts of sublethal mercury exposure on birds: a detailed review. *Rev. Environ. Contam. Toxicol.* 244, 113–163.
- Williams, M., Todd, G.D., Roney, N., et al., 2012. Toxicological Profile for Manganese. Agency for Toxic Substances and Disease Registry (US), Atlanta (GA).